
UNM PEDIATRIC ED SEXUAL ASSAULT & NEEDLESTICK GUIDELINE nPEP GUIDELINE PAGE 2



^{*} Based on event location jurisdiction.

Created May 2025

Contributors: Rebecca Girardet, Martha Muller, Jeremy Snyder, Pree Sarangarm, Jacob Kesner, Tara Neubrand, Yomayra Perez Morel, Kenny Wilson, Sara Skarbek-Borowska, and David Jolley

[†] Urine, vaginal, rectal, or pharyngeal depending on sites of contact (self-swab if age-appropriate)

[§] Patients with female reproductive organs only

HIV Post-Exposure Prophylaxis

National Clinician Post Exposure Prophylaxis PEPline (888) 448-4911

High Risk • Exposure of: Mucous membranes, non-intact skin, percutaneous contact • With: Blood, semen, vaginal secretions, rectal secretions, breast milk, or any bodily fluid visibly contaminated with blood - 72 hrs since exposure - 272 hrs since exposure - 72 hrs since exposure - 100 Risk • Exposure of: Mucous membranes, non-intact skin, percutaneous contact • With: Urine, nasal secretions, saliva, sweat, tears - 100 Risk • Exposure of: Mucous membranes, non-intact skin, percutaneous contact • With: Urine, nasal secretions, saliva, sweat, tears

So you've decided to start nPEP Age < 13 Provide 28 Days Supply: Baseline labs: **STOP** Call Peds ID CBC, Diff, Cr, AST, ALT Raltegravir: • 3 - < 6 kg: 25 mg BID • 6 - < 10 kg: 50 mg BID • 10 - < 14 kg: 75 mg BID No • 14 - < 20 kg: 100 mg BID Can the patient swallow pills? 20 - < 28 kg: 150 mg BID • 28 - < 40 kg: 200 mg BID • ≥ 40 kg: 300 mg BID Yes 25 & 100 mg are chewable and 100 mg can be split. The 25 mg tabs can be crushed and dissolved in 5 mL of liquid No ≥ 25 kg? Lamivudine (10 mg/mL): 5 mg/kg BID (max 150 mg) Yes Zidovudine (10 mg/mL): • 4 - < 9 kg: 12 mg/kg BID Biktarvy 1 tab Qday 28 days • 9 - < 30 kg: 9 mg/kg BID Bictegravir/emtricitabine/tenofovir ≥ 30 kg: 300 mg BID

nPEP Follow-up:

Age 18:

- Send a Tiger Connect Message to Truman Clinic HIV Connect to Care Line Team to schedule appt for follow-up within 7 days
- Give adolescent nPEP DC instructions under PEM Clinical Resources on Smartsheets.

Age 13-17:

- Follow-up with Para Los Ninos within 7 days
- Give child nPEP DC instructions under PEM Clinical Resources on Smartsheets

Age < 13:

Discuss nPEP follow-up with Peds ID Attending

If nPEP prescribed, please call the receiving pharmacy (if open) to be sure it is in stock and covered by patient's insurance. This is why **UNM Discharge Pharmacy is preferred**. Pharmacies below.

Needlestick: Follow-up with Peds ID if starting nPEP

Pharmacies with nPEP medications:

UNM Discharge Pharmacy (PREFERRED) Can only fill if > 2 hrs prior to closing time and dc order in Meds can be delivered to the Peds ED before patient leaves	2211 Lomas Blvd NE, Albuquerque NM 871206	M-F 8A-7P, Sat 8A-5P, Sun 9A-4P	(505) 272-1591
Truman Health Services Pharmacy (UNM Medical Group, Inc)	801 Encino Pl NE, Ste B4, Albuquerque NM 87102	M-W 8:30A-5:30P, H 8:30A-7P, F 8:30A-5:30P)	(505) 925-0996 After Hours (833) 440-1144
Southwest Care Jefferson Clinic and Pharmacy	4710 Jefferson Street NE, Suita A	M-F 8:30A-5:30P with break Wed 1P-2:30P	(505) 780-4044
Walgreens #4649 (Raltegravir may not be ready for next day pickup)	3501 Lomas Blvd NE, Albuquerque NM 87106	M-F 9A-7P, Sat 9A-6P	(505) 255-8908

References:

AIDS Education & Training Center. nPEP Quick Guide for Providers. National AETC Support Center. Nov 22, 2021. https://aidsetc.org/resource/npep-quick-guide-providers

Bechtel K, Bennett BL. Evaluation of sexual abuse in children and adolescents. Post TW, ed. UpToDate. UpToDate Inc. Accessed May 5, 2025. https://www.uptodate.com/contents/evaluation-of-sexual-abuse-in-children-and-adolescents

Buchanan JA. Evaluation and management of adult and adolescent sexual assault victims in the emergency department. Post TW, ed. UpToDate. UpToDate Inc. Accessed May 5, 2025.

https://www.uptodate.com/contents/evaluation-and-management-of-adult-and-adolescent-sexual-assault-victims-in-the-emergency-department

Children's Hospital of Colorado Clinical Pathways and Measures Committee. Blood or Bodily Fluid Exposure in the Community (Non-Occupational). In: *Clinical Pathway's*. Children's Hospital of Colorado, July 24, 2024.

https://www.childrenscolorado.org/globalassets/healthcare-professionals/clinical-pathways/blood-or-bodily-fluid-exposure-in-the-community-non-occupational external.pdf

Jackson AM, Artis AR, Atkinson NK. Adolescent sexual assault. In: Kliegman RM, St. Geme JW III, Blum NJ, et al., eds. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:(Ch) 162.

Kent CK, Casey CG, Dott M, et al. Sexual assault and abuse and STIs – adolescents and adults. In: *Sexually Transmitted Infections Treatment Guidelines, 2021*. The Center for Disease Control and Prevention; July 22, 2021. Accessed May 5, 2025.

https://www.cdc.gov/std/treatment-guidelines/sexual-assault-adults.htm

Molnar J, O'Connell MS, Mollen C, Scribano P. Sexual assault: child and adolescent. In: Shaw KN, Bachur RG, Chamberlain J, Lavelle J, Nagler J, Shook JE, eds. Fleisher & Ludwig's Textbook of Pediatric Emergency Medicine. 8th ed. Wolters Kluwer; 2020:(Ch) 127.

Schillie S, Vellozzi C, Reingold A, et al. Prevention of hepatitis B virus infection in the United States: recommendations of the advisory committee on immunization practices. In: *Morbidity and Mortality Weekly Report (MMWR)*. The Center for Disease Control and Prevention; Jan 12, 2018. https://www.cdc.gov/mmwr/volumes/67/rr/rr6701a1.htm

Workowski KA, Bachmann LH, Chan PA, et al. Sexually transmitted infections treatment guidelines, 2021. In: *MMWR Recommendations and reports: Morbidity and mortality weekly report Recommendations and reports.* The Center for Disease Control and Prevention; July 23, 2021. https://www.cdc.gov/std/treatment-guidelines/STI-Guidelines-2021.pdf