



# SCHOOL OF MEDICINE

DEPARTMENT OF  
ORTHOPAEDICS & REHABILITATION

**Gehron Treme MD, Andrew Veitch MD, Dustin Richter MD, Christopher Shultz MD,  
UNM SPORTS MEDICINE**

1213 University Blvd NE, MSC10 5600, 1 University of New Mexico; Albuquerque, NM 87131-0001

Phone: 505-272-2231 Fax: 505-925-4015

## **Rehabilitation Protocol**

### **Achilles Tendon Repair and Non-operative Protocol**

#### **Phase I**

2 weeks post op/post injury

- Start physical therapy
- Walking boot with heel lift insert – Weight bearing as tolerated
- Inversion/eversion exercises—iskinetics, tubing, manual resistance, etc.
- Active dorsiflexion exercises to increase ROM
- Passive dorsiflexion to neutral only
- Modalities as needed
- Warm whirlpool
- No resisted plantar flexion (active and passive plantar flexion motion ok)
- Talocrural joint mobs as needed

#### **Phase II**

6 weeks post op/post injury

- Remove walking boot - WBAT
- PRE's for gastroc/soleus strengthening as tolerated (prevent Achilles tendonitis)
- Work to increase full gastroc/soleus ROM
- Proprioception exercises—single leg stands, stable and unstable surfaces
- Normalize gait

#### **Phase III**

12-16 weeks post op/post injury

- Full ROM
- Normal gait
- Near full strength—able to stand on toes of affected side in full plantar flexion, or able to do 4x25 single foot heel raises
- Start jogging program if strength acceptable, no limp with gait, and ROM is full